

# REPORT DOCUMENTATION PAGE

Form Approved  
OMB No. 0704-0188

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503.

1. AGENCY USE ONLY (Leave blank)		2. REPORT DATE		3. REPORT TYPE AND DATES COVERED FINAL 01 Apr 97 To 31 Dec 97	
4. TITLE AND SUBTITLE Nd:YLF Laser System for the Study of Termolecular Reactions and Metastable Isomers of Polyatomic Molecules				5. FUNDING NUMBERS F49620-97-1-0255 2303/ES 61102F	
6. AUTHOR(S) DR ROBERT E. CONTINETTI					
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) University of California, San Diego Department of Chemistry - 0314 9500 Gilman Drive La Jolla CA 92093-0314				AFRL-SR-BL-TR-98-0316	
9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES) AFOSR/NL 110 Duncan Ave Room B115 Bolling AFB DC 20332-8050  Dr Micahel R. Berman				10. SPONSORING / MONITORING AGENCY REPORT NUMBER	
11. SUPPLEMENTARY NOTES					
12a. DISTRIBUTION / AVAILABILITY STATEMENT  Approved for public release; distribution unlimited.				12b. DISTRIBUTION CODE	
13. ABSTRACT (Maximum 200 words) The DURIP grant F49620-97-1-0255 has supported acquisition of a state-of-the-art regeneratively amplified Ti:Sapphire laser system. This laser system is the primary laser for our new multi-particle photoelectron-photofragment coincidence spectrometer, and is currently being used to carry out experiments on three-body dissociation dynamics in clusters of oxygen and ozone. These experiments make use of a new particle detector built in our laboratory capable of recording the time- and position-of-arrival of up to eight photofragments in a period of 20-500 ns following a laser pulse. The initial DURIP proposal discussed the purchase of a Nd:YLF psec regenerative amplifier, however, the Ti:Sapphire laser purchased is a much more flexible and powerful laser, producing both psec and fsec pulses with a higher peak power. The initial experiments with this laser will use the fundamental and the 2nd, 3rd and 4th harmonics. Future applications will include the generation of tunable radiation with an optical parametric amplifier and ultrafast experiments using 100 fs pulses. We took delivery of the laser December 1, 1997, with initial installation December 12, 1997.					
14. SUBJECT TERMS				15. NUMBER OF PAGES	
				16. PRICE CODE	
17. SECURITY CLASSIFICATION OF REPORT (U)	18. SECURITY CLASSIFICATION OF THIS PAGE (U)	19. SECURITY CLASSIFICATION OF ABSTRACT (U)	20. LIMITATION OF ABSTRACT (UL)		

19980421 059

**Final Technical Report - AFOSR DURIP Grant #F49620-97-1-0255**

**4/1/97 - 12/31/97**

**Nd:YLF Laser System for the Study of Termolecular Reactions and Metastable  
Isomers of Polyatomic Molecules**

**Principal Investigator -- Robert E. Continetti**

**University of California, San Diego  
Department of Chemistry - 0314  
9500 Gilman Drive  
La Jolla, CA 92093-0314**

**Tel: (619)-534-5559  
FAX: (619)-534-7042  
email: rcontinetti@ucsd.edu**

## 1. Equipment Acquired under DURIP Grant

The DURIP grant F49620-97-1-0255 has supported acquisition of a state-of-the-art regeneratively amplified Ti:Sapphire laser system. This laser system is the primary laser for our new multi-particle photoelectron-photofragment coincidence spectrometer, and is currently being used to carry out experiments on three-body dissociation dynamics in clusters of oxygen and ozone. These experiments make use of a new particle detector built in our laboratory capable of recording the time- and position-of-arrival of up to eight photofragments in a period of 20-500 ns following a laser pulse. The initial DURIP proposal discussed the purchase of a Nd:YLF psec regenerative amplifier, however, the Ti:Sapphire laser purchased is a much more flexible and powerful laser, producing both psec and fsec pulses with a higher peak power. The initial experiments with this laser will use the fundamental and the 2nd, 3rd and 4th harmonics. Future applications will include the generation of tunable radiation with an optical parametric amplifier and ultrafast experiments using 100 fs pulses. We took delivery of the laser December 1, 1997, with initial installation December 12, 1997.

**Acquired Equipment :** CPA-2000 kHz Ti:Sapphire regenerative amplifier,  
producing either 0.4 W, 1.4 ps pulses at 1 kHz or  
0.8 W, 120 fs pulses at 775 nm

**Manufacturer :** Clark-MXR, Inc.

<b>Cost (Including CA State Tax and Shipping):</b>	\$185,653
DURIP Funds	\$122,268
UCSD Matching Funds and other non-Federal Funds	\$ 60,222

## 2. Research Projects

The new photoelectron-multiple-photofragment coincidence spectrometer in our laboratory allows a new generation of experiments on three-body dissociation dynamics to be carried out. The CPA-2000, with its picosecond and femtosecond pulse capabilities, plays an integral role in these new experiments. We plan on applying this apparatus to a study of some of the three-body association reactions important in atmospheric and combustion phenomena, such as the recombination reaction that produces ozone:  $O + O_2 + M \rightarrow O_3 + M$ . In the initial experiments using the new laser, the dissociative photodetachment and photodissociation of  $O_4^-$  and  $O_6^-$  have been studied at 775 nm and 388 nm. The results at 775 nm show that while  $O_4^-$  only undergoes dissociative photodetachment at that wavelength,  $O_6^-$  only exhibits photodissociation processes yielding  $O_2^- + O_2 + O_2$ . This suggests that either a breaking of the high symmetry of the  $O_4^-$  core or the presence of a new charge-transfer-to-solvent band in  $O_6^-$  has a profound impact on the photochemistry at this long wavelength. At 388 nm, the results confirm our previous observations at 532 and 266 nm that while the addition of  $O_2$  to  $O_4^-$  to form  $O_6^-$  only slightly perturbs the dynamics of dissociative photodetachment, significant changes in

the photodissociation dynamics occur, yielding highly vibrational excited  $O_2^-$  products. We are now beginning our studies of the  $O_3^- \cdot H_2O$  system, which will provide a model for the ozone recombination reaction.